


# Dermatology



- Reduction of subcutaneous fat by topical medication of lipolytic agents
- Reduction of subcutaneous fats by topical application of lipolytic agents
- Risk factors for lipodystrophy in HIV patients considering genetic factors
- Comparison of histochemical staining in malign melanoma – which staining is best-suited?
- Diagnostic validity of microscopic grey scale analysis in different skin tumors
- Determinants of survival in malignant melanoma – a 10-year analysis
- Sensitivity and specificity of the dermatoscopy in melanocytic nevi and malign melanoma
- Comparison of prognostic validity of several tumor markers in malign melanoma
- Predictive value of the Su-Par in malign melanoma – a model development by combining prognostic parameters
- Outcome in basal cell carcinoma – a multicenter study with 2000 patients
- Incidence of malign melanoma between 1979 and 2003. Description of modification of localization, histological subtypes and patients' characteristics
- Sensitivity and specificity of antibodies in dermatomyositis and polymyositis
- Evidence of auto-antibodies in patients with collagenase and non-rheumatic inflammatory diseases
- Diagnostics and medical treatment of drug induced exanthemas
- Success rates of hyposensibilisation in patients with anaphylactic reactions after wasp stings
- Comparison of different treatment procedures of Osler's disease
- Prospective studies in patients with lipoedema and liposuction treatment
- Lymph therapists' standard of knowledge regarding diagnostics and therapy of lipoedema
- Comparison of treatments of lipoedema in different professional Groups
- Comparison of lipoedema treatments in different occupational groups
- 6 month-results after liposuction in female patients suffering from lipoedema
- Long-term outcome after liposuction in patients with lipoedema
- Definition of new measuring methods to determine body measurements of female patients suffering from with lipoedema
- Quality of life after liposuction in female patients with lipoedema
- Prevalence and clinical symptoms of the lipoedema in GP care
- Venous and lymphological diseases of the lower extremities in women – an epidemiologic survey including 1000 women
- Coriumdicke vor und nach KPE bei Patientinnen mit Lymphödem
- Prevalence of varicosis in GP care – evaluation of different therapeutic regimes